

THE EISENHOWER LIBRARY



3 1151 02718 8360

123558


Library



Johns Hopkins University









THE GREEK MUSEO
AND THE RECONSTRUCTION OF THE
IN THE PERSIAN, SIGACONIA LICHILAR.

BY
EDWARD WILLIS CROWE.

DISSERTATION
SUBMITTED TO THE BOARD OF UNIVERSITY STUDIES OF THE
JOHNS HOPKINS UNIVERSITY
IN CONFORMANCE WITH THE REQUIREMENTS FOR THE DEGREE OF
DOCTOR OF PHILOSOPHY,
BALTIMORE.

1908.

INTRODUCTION.

Through the kindness of Professor W. W. Brown, it was made possible for me to go to Beaufort, N. C., in the summer of 1932, and while there, I began, at his suggestion, to collect material for the development of the head-skeleton of the Pipefish. I soon found young embryos and segmenting eggs, and, wishing to take up the embryology of this fish, I deferred the former work until a later date.

The collection of further material and the observations on the breeding habits were made at Beaufort, during the summers of 1933 and 1934; also, with running sea-water as here, the difficulties necessarily attendant on this work were materially reduced.

This preliminary work was done in the laboratory of the U. S. Bureau of Fisheries at Beaufort, N. C. I am indebted to the Commissioner, Hon. George W. Brown for the opportunity to make use of the most excellent facilities at hand there. To the Director, Dr. Cassell Trave, I am under obligations for many helpful suggestions.

The further work was done in the Biological Laboratory

of the Johns Hopkins University. To Professor W. L. Brown, I am very grateful for the interest taken in my work and for advice and direction. I also wish to thank Mr. F. A. Andrews and Mr. Caswell Crane for advice in overcoming the technical difficulties of my work.

MATERIAL AND METHODS.

Male Pipefishes with full pouches were drawn into the laboratory and there the upper end of the pouch was opened with forceps and a finger removed and put under the microscope. If these were in a stage where the head of the fish was cut off, the flaps of the pouch still open with scissors and removed (frequently bringing eggs with them), the eggs removed by tracing with needles the connective tissue binding them down. If the eggs were too young, the fish was put back in running water and examined again later; although they rarely survived a second operation, unless the eggs were nearly hatched and hence came out easily. This is a wasteful process since many eggs are spoiled in removing them. The obtaining of a series of eggs and embryos in Siphostoma, is a long, slow, and laborious task, and is quite as much the result of chance as of skill and knowledge.

A variety of killing fluids has been used. The oil drops under the germ disc were so thickened by osmic acid and Flemming's fluid, that these reagents could not be used. Acetic alcohol, Heinenberg, sublimate-acetic, micro-acetic, all gave good blastodermis but the yolks generally went to pieces. Excellent results were obtained with fresh Formalin, 1% and 2% formalin, and, for later stages, Wilson's and Worcester's fluids. This latter is one of the best fluids for killing teleostegs with which I am acquainted. It is composed of saturated solution in 10% formalin, 10 parts, phenyl acetic 10 parts. The eggs were left in this from 30-60 minutes, washed 4 times, run up into 70% alcohol and the excess of alcohol removed with 100%.

The eggs, placed in an embryo under cover from the water killing fluids, were run through into a 10% solution of hypochlorite of Sodium for 24 hours to a firm translucent blastoderm the transparent of chorion. Good success with these fluids was very sensitive to the blastodermis, and generally the eggs were run up into 70% alcohol and the excess removed with 100%.

The younger blastodermis were picked off the yolks

and efficiency, but the probability is increased from the point of view of the insecticide to get the maximum effect. The first stage was to use these eggs were not about, the for this purpose those killed in Varenzi's fluid, on account of their soft yells, were especially good. The point of view killed in Varenzi, if they had been in the, then to remove them, would be much more rapid than the other as quickly as possible.

In order to obtain which eggs in the Varenzi, it is necessary to start them. By putting them in this strength during the time for from 30 to 45 minutes, the embryonic tissues have the ability to the yells, and there is a real resistance on yells.

The eggs were incubated in paraffin, and a distance of from 5 to 10 microns from the surface either in Varenzi's fluid or in the fluid of the Varenzi. The former gave the best results and was so easy to handle that it was almost exclusively used.

HABITAT.

Pipefishes are found in all the warm and temperate regions of the world, but are not exclusively marine. DeMe (1935) reports that C. macrostichus was found in the Ganges-China Hill's above tide limits. Arnold (1938), he finds that C. sticticus, latipectus, carce and three species of Polyichthys go up the rivers of India. Sanchez ('04) reports Polyichthys cauca and fluvialis in the rivers of the Ibero-Peninsula.

In the harbor at Newport, in quiet shallow waters where there are sandy bottoms, masses of Ulva abound and in one of the Pipefishes live. On fishing in nets with a fine-meshed seine, they may be caught in considerable numbers.

It may be well to note that the color of these fishes changes with the sea-weeds upon which they are found. C. florida among tufts of sandy sea-weed is dark-green, but put into aquaria with Ulva or Cladophora, it becomes bright green. C. fuscus is ordinarily of a dull brown color, but several specimens caught in a tide pool filled with red sea-weed were brick-red in color and from this were thought to be a new species.



other, individuals touching and the male fish, the male
more demonstrative. Just before the actual transfer, the
male becomes violently excited and demonstrative, and
the male and anterior body-parts in a compressed fashion
and with his snout crosses the female on the left. The
female responds to this but does not become so excited.
This is repeated several times, and female feeding more
excited and time they touch each other. In a very quick
as a flash, the sexual embrace takes place and then the
fishes separate to begin again in a few minutes.

This embrace consists in the fishes interlocking their
bodies like two capital letter S's, the one reversed
on the other thus bringing their face to face. Thus they
hold their bodies together while the eggs pass from the
oviduct into the pouch. Their bodies touch in three places;
in the anterior region, just back of the pectorals;
in the posterior region, at a point about two-thirds of
the way from the anus to the caudal; and at the anal open-
ings. The anal ptyllia, or the protruding oviduct of the
female, is, at the moment of contact of their bodies,
insert into the button-hole-shaped opening at the anterior
end of the marsupium. The eggs, in number a dozen or more,
now pass into the pouch and are presumably fertilized by this

movement.

The legs are now in the anterior end of the pouch, and no more can be received until they have been shifted into the posterior end. To bring this about, the male performs some very curious movements. He stands nearly vertically, and, resting his caudal and a small part of the tail on the floor of the aquarium, bends backward and forward and twists his body spirally from above downward. This is repeated until the legs have gotten down into the posterior end of the pouch. I do not think that any other means than the above are used to bring this about. The pouch in a "pithon" fish was opened and examined scattered over its inner surface, but there was no evidence of ciliary action. Sections from pieces of both dorsal and ventral parts of the sac killed in formalin, fixative fluids (Worcester), failed to show cilia.

Then for some time the animals remain quiescent, the male assuming the form of a broad first capital U. The head is extended in a nearly horizontal direction, and the body in the region of the middle of the tail touches the floor of the aquarium. This position is retained for a time varying from five to ten minutes. Convulsive movements, lasting only for a moment, may take place.



The processes above described are repeated until the pouch is full. In one pair the first copulation began at 8:45, and the second at 17:30. In another pair there were four contacts as follows, 10:10, 10:34, 10:33, at which time the males were only half way down the pouch; and at 11:00. These observations were made at night, between 8:45 and 11:30, in the brightly lighted laboratory. It is very probable that the transfer takes place at any and all hours of the night. It may be noted in passing that the fishes seemed entirely unaffected by the lights. No attempt was made to determine the sex of Lafont.

It does not seem likely that all the eggs are transferred at once; first, because of the curious means used to move them backwards in the pouch, secondly, because males are frequently found with the pouch only half filled, thirdly, because males with full pouches are not infrequent. These processes have been repeated several times the animals are seemingly exhausted and remain quiet for at least two and one-half hours (the extent of my observations). On this same night, a third small male in an aquarium with three females "courted" two of them alternately, but no

transfer was made. These last had protruding oviducts. For coition to take place, it seems necessary that the fishes should be nearly equal in size. I raise female, priced with a single three-fifths per size, through sex pots into the water.

It is common to observe a series of two rows of eggs parallel in other lower vertebrates. Jordan ('31) records for *Misgonyx* a very interesting series of observations of a courtship, lasting several hours, in which the male displays his account of the female's place in important ways. Near ('0), in the spawning of *Lepidosteus*, described how the males with wide-spread fins, swim around the females and press their snouts together. He is sure a courtship unknown among the Invertebrates. Ravitza ('04) has described how the male of *Octopus vulgaris* strokes the processes of the female. All these contacts seem to be intended to excite the animals preparatory to the sexual act.

The arrangement of eggs in the pouch depends wholly on the size of the latter. There are always two sets of eggs, one on each side. Each set may consist of one, or two, or of three rows of eggs, and these may be one or two eggs deep. As noted there may be one, two, or even three deposits of eggs into one pouch. In what order these

young would escape from the pouch, I was not sure. Un-
 fortunately the boys were so glib that they did not
 do so from the young.

The time at which hatching can be given to the young
 (with a variation of light hours) after one fertilized egg.
 These young live four days, feeding on yolk, with the
 semi-circular motion of the head, and the same semi-circular
 mouth called found in the parents. It explains that, when
 the father dies four days after the hatching, the little
 fishes were with fresh shells.

The eggs within twenty-four hours after deposition
 can easily be extracted from the pouch, coming out in
 masses, without injuring the father. In two cases, males
 relieved of eggs, received a fresh incubation the following
 night. One of these, stripped the second time, died
 after taking on a third lot. When the male have been
 in the pouch thirty-six or forty-eight hours, they become
 finally fastened to it both at top and bottom, so that it be-
 comes necessary to kill the fish and then cut away the flaps
 of skin before one can extract the eggs.

The fishes vary in size. The extremes in egg-bearing
 males of S. floridae I have found to be 4.5 to 6.9 inches, and
 in females 3 to 8.4 inches. As a general rule however the
 females are somewhat the larger.

Foot-note to p. 11.

A review of the literature, from the time of Aristotle to the present, on the sexual characters, breeding habits, and embryonic structures of the lampbrushes, together with certain observations of my own, is now ready for publication.

THE SECRETION OF THE EGG IN THE TELEOST --
SIPHOSTOMA FLORIDANUM.

I. THE OVARIAN EGG.

The ripe egg of this fish is of fairly good size having a diameter of about 1 millimeter. It possesses a thin transparent membrane or shell, which, under the one-twelfth homogeneous oil immersion lens, shows no structure in sections, but in surface views presents, when stained lightly with Madderine, a notably punctate appearance. These membranes were generally removed after killing the eggs, but if left on the eggs do not get very hard and offer no obstruction to imbedding and sectioning processes. The eggs are formed in ovaries which, viewed from without, present the ordinary Y shaped structure common to the Teleosts. These ovaries are two tubular organs situated in the posterior dorsal portion of the body cavity, and are confluent behind to form the short oviduct which opens in the posterior lip of the anal aperture.

However when one of the ovaries is sectioned, a very interesting structure is revealed. Running lengthwise throughout the whole extent of the ovary is a raphe situated about two-thirds of the distance from one wall. From this raphe are budded off in succession to form a spiral of eggs which surrounds the raphe, the outermost egg being the oldest and largest. As this egg ripens, it markedly increases in size and crowds the other eggs together with the raphe closely to one side of the tube. In the ovaries of older and larger fishes, two or three eggs may ripen side by side and then the raphe and its young eggs are very much crowded and contorted. As the eggs become ripe they enormously distend the ovaries both in diameter and length -- in length until they frequently extend forward to the region of the stomach. At this time females ready to spawn are noticeable for their greatly distended abdomens.

The young eggs, as first pointed out by Cuvier (1837) have large nuclei with several nucleoli, but in the older ovarian eggs the germinal vesicle is not so apparent. The grown eggs, still attached in the ovary are surrounded by a layer of peripheral oil drops. This same structure persists in the eggs after extrusion, so that the germinal vesicle can not be seen. The sections I have made of eggs just extruded

we so far as possible have followed the sections we figured in our paper. As a result, the older observers, Rastus ('53), Rastus ('55, '57, '60), Vogt and Pappenheim ('59), although they studied the ovary with the microscope, missed these peculiar structures. Later observers, Sobel, Cohen, and others ('57), and Hunt ('62) have made sections but have not gone very far into the structure, nor will I myself do so now, since it is my intention to work up the structure and development of this organ later, the material for this being now on hand.

II. THE MANTLE OR PROTECTION.

This has already been described in the first part of this paper, but it will be well to emphasize the fact that the process is such as to prevent absolutely any contact of the eggs and sperms with the sea water.

III. FERTILIZATION.

The egg of *Cirrhosoma floridanus*, as before mentioned, possesses a very thin or perfectly transparent shell. This surrounds an egg made up of straw colored yolk having many orange-red oil globules imbedded in its periphery, and these surrounded in turn by a thin pellicle of protoplasm. The colored oil globules render the egg so perfectly opaque that

I have never been able to find the micropyle.

It is strange to say, the egg of a related European form, Glyptothorax ocellatus, was the first fish and possibly the first vertebrate egg in which this opening was discovered. Whether this egg is transparent or not I can not say, but in it Doyère ('49) found the micropyle just over the "disque proli-gero" and gave its diameter as 1/110 mm.

A. Natural Fertilization.

Different investigators vary in their conclusions, or more correctly, their conjectures, as to the time of fertilization. A priori, one would expect the fertilization to be effected at the time of transfer. Probably the surest way to determine the time of impregnation would be to take a male immediately after the transfer, cut through the pouch just back of the forward end behind the genital opening, and then examine the eggs in the hinder part of the pouch for spermatozoa. This I had intended to do during each of the past summers. Although I had numerous transfers between fish kept in aquaria each summer, yet I saw the copulation on one night only (in 1933.) between two pairs of fish. The seeming absolute necessity for keeping these fish for the early stages of segmentation, prevented my sacrificing either to determine this point.

Suol ('32), Koster ('32), Liljebom ('31), and others, think that the fertilization takes place at the time of copulation, while A. W. Salm ('34), and Groves ('35) think that it follows later, and Ekstroem ('31) believes it takes place while the eggs are in the pouch. My own belief is that sperms and ova are emitted simultaneously and while I have no direct evidence, I wish to adduce the following facts corroboratory of this conclusion.

I believe that the extraordinary "siesta stiel", or period of sexual excitation of these fishes, described above, is intended to prepare them for the mutual discharge of the sexual products. In the description of the copulation and attendant phenomena I have called attention to similar sexual excitements in Amphibia, Cephalopoda and Chytridophora, which are preparatory to the discharge of sperms as well as of eggs.

But the second set of phenomena is still more strongly corroboratory. On July 8, 1924, two fish were paired and during the night mated copulatively. They remained in the same aquarium for four days, and then the female was killed, her ovaries excised, cut up, and put into fixing fluids while some of the ovarian eggs, which fell into the body cavity, were also killed. When these eggs were examined some months later, among them were found two embryos with the blastopore closed. None of the other eggs showed any trace whatever

of impregnation. Again two lots of eggs from pouches killed in 1892 were examined two years later and found to be in the eight to sixteen-celled stage. In one lot, however, there was found an embryo with black eyes and long tail, and in the other two eggs in which the blastodermis covered one-half, the embryos one-fourth, of the circumference of the egg. These two lots of eggs had never been removed from the salls, and these salls were still bound together in masses as they came from the pouch. Thus all chance of the eggs having been mixed is eliminated. Again a lot of eggs put up in August, 1904 were found to be in the eight-celled stage, but among them were found two embryos with pectoral fins.

It is true that in opposite ends of the pouch eggs of different layings and consequently ^{of} different ages are found, but never with differences of age more than thirty-six hours against about three to five days in the above cases. From these facts I can draw but one conclusion, that at the time of coition both spermatozoa and ova are simultaneously extruded, and, as the female withdraws her oviduct from the buttonhole-shaped opening of the marsupium, sperm lodge on it and work their way through it into the ovary and there fertilize eggs. This happens only occasionally, but it seems to me a strong proof of my contention as to the time of fertilization.

U. Artificial Fertilization.

twice

This was tried by the wet method and once by the dry. The eggs and the torn up testes were thoroughly mixed in sea water, and after a few minutes were aerated in stirred sea water. From a third lot of eggs, the water was carefully drained, and over them was poured sperm from testes which had been torn up in a perfectly dry dish. These were allowed to stand for a few minutes and were then placed in clean running sea water. The females were certainly ripe for spawning, and the males were well grown and had not recently borne eggs, so they were presumably fertile. A control experiment was made by putting a batch of this last lot in running sea water without the addition of sperm. In all cases the results were the same. At the end of one and one-half hours, protoplasm could be seen collecting at the upper pole. After two to three hours, it was noticed that the eggs had flattened slightly at the animal pole and that there was being formed a pretty clearly defined round germinal disc resting on a layer of orange-red oil drops. At an age of four to six hours, the germinal disc was at its prime, but neither then nor at any subsequent time was there any trace of segmentation. From this time on, the germinal disc gradually lost its sharp outlines, flattened down, and went

to pieces. In one lot of eggs at the age of twenty-six hours, the germinal disc had gone bad; in another after twenty-five hours, it was no longer round and its edges were irregular and fragmentary; in a third lot less than ten percent of the eggs, were alive after twenty-three and one-half hours.

These eggs were all alike save that in one lot, some when taken from the ovary, showed a very faint aggregation of protoplasm at the germinal pole, while in another lot ^{latter condition} the eggs were of unequal size. This, is, however, by no means an uncommon occurrence. Such differences are met with repeatedly in my preserve material, where eggs one-half to two-thirds the size of the normal ones are found. Save that the blastodermis are somewhat smaller, there is nothing unusual about the development of these small eggs. In this connection, Brock ('06) says that the eggs of the Herring vary in size in the same fish or in fishes of different localities, but thinks that this in no wise affects their development.

From my experiments it seems pretty clear that artificial fertilization is not possible in the Pipefish, thus confirming the a priori opinion that this would not take place in fishes provided with such extraordinary apparatuses for the deposition and impregnation of the eggs, without their ever coming in contact with the water. Since the eggs will

the first part of the paper, we consider the case of a single agent.

In the second part, we consider the case of multiple agents.

In the third part, we consider the case of a single agent.

In the fourth part, we consider the case of multiple agents.

In the fifth part, we consider the case of a single agent.

In the sixth part, we consider the case of multiple agents.

In the seventh part, we consider the case of a single agent.

In the eighth part, we consider the case of multiple agents.

In the ninth part, we consider the case of a single agent.

In the tenth part, we consider the case of multiple agents.

In the eleventh part, we consider the case of a single agent.

In the twelfth part, we consider the case of multiple agents.

In the thirteenth part, we consider the case of a single agent.

In the fourteenth part, we consider the case of multiple agents.

In the fifteenth part, we consider the case of a single agent.

In the sixteenth part, we consider the case of multiple agents.

In the seventeenth part, we consider the case of a single agent.

In the eighteenth part, we consider the case of multiple agents.

In the nineteenth part, we consider the case of a single agent.

In the twentieth part, we consider the case of multiple agents.

In the twenty-first part, we consider the case of a single agent.

In the twenty-second part, we consider the case of multiple agents.

In the twenty-third part, we consider the case of a single agent.

live for some twenty hours in sea water, it must be due to spermatozoa which are histologically affected by it. It has long been known that the sperms of both salt and fresh water fishes lose their vitality if left in the water any time and can not impregnate eggs. Quatrefages first ascertained this for the Pike and other fresh water fishes. Hoffmann ('01) says that the sperms of *Scorpaena* die quickly in salt water. Reichard ('03) found that the sperms of the "Hill-Fish" Pike die after one minute in the water.

In this connection the experiments of Huot ('02) are very interesting. He took the eggs of *Syngnathus aequilii* from the marsupium of the male, and ^{being} careful not to break the egg membranes (these eggs were presumably fertilized), put them in clean aerated sea water. This he did also with eggs just before deposition (ovarian eggs), but in no case did development go on more than a few hours. When he put into the water larvae old enough to move freely, but these too died within forty-eight hours. I can confirm all his results. I have found that eggs in segmentation will go on dividing for a short while, but that within eighteen hours all die. The discoveries of Huot ('02) and ^{of} John ('04), that the pouch and its contents act as a physiological placenta, offer the explanation for the above phenomena. The eggs and embryos,

depending on this for oxygen and food, can not exist out of the pouch.

IV. MATURATION.

Unable to fertilize artificially the eggs of Sinuostoma floridae, and having found it impossible to get from the pouch eggs young enough to show the formation of polar bodies, I am unfortunately not in position to say anything of the process of maturation. For the latest and best work on this phenomenon the reader is referred to Pearson's paper ('36).

V. FORMATION OF THE GERM DISC.

In the Pipefish, fertilization is not necessary to bring about the formation of the germinal disc. Immersion in water supplies the stimulus as it does in many other fishes. All workers on the Salmonids, Ziegler ('02), Wis ('33), and others, so report. Kowalewski ('06) found it true for the Goldfish, as did Agassiz and Whitman ('55) for Otenoldubrus, though they state that for pelagic eggs the germ disc is generally not formed until after impregnation. Prook ('36) confirms this for the Herring, but I have found that the eggs of the Sargassumfish, Pterophryne histrio, form the germ disc shortly after extrusion. Hertwig says (Mambuch p. 544), "One can emphatically say for almost all fish eggs that, by

their transfer into water, such a powerful force is brought into play that the concentration of the germ disc results", but that "if they are impregnated first, a more rapid growth and larger size for the germ disc follows".

All writers, notably Brook ('38) and Ender ('57), describe this formation as brought about by the streaming of the protoplasm to the germinal pole. There are three modes in which this may take place:

I. By streams from the circumference only. This is the method in most fishes, especially those with pelagic eggs. See Brook, Aydar, Kingsley and Conn, and many others.

II. By streams from the circumference with the help of little "processions" from the interior of the yolk (Ziegler ('32) and Oellacher ('72) for the crust).

III. In all directions from the yolk, the streaming goes to the germinal disc (Carassius, Kowalewski ('36)). As best I can determine, the Pipefish comes under class two. This matter will be further referred to in the section dealing with the periblast.

Intimately connected with the foregoing is the collecting of the oil drops underneath the germ disc. In pelagic eggs, the oil is in one great globule generally near the

center of the yolk, but in the Pipefish many small orange-red globules are imbedded in the periphery of the yolk. When the protoplasm moves up to the animal pole, the oil globules go also and are collected under the germ disc to form the "disque huileux" of Lereboullet. This is a phenomenon very common among Teleosts. It has been reported by all workers on the Salmonoids, by Hanson ('37) for the Stickleback, Kowalewski ('80) for Carassius, and by many others. Rathke ('37) first described these processes in Pipefishes from the Black Sea. He says that the germinal disc is formed after the eggs come into water, and that the yellow-red "fat" drops which surround the yolk flow up to and spread out under the disc in a layer covering about one-third of its upper surface. Kupffer ('68), describing the egg of a European Syngnathus, says, "This fat forms a mass of drops of different sizes which encloses the germ disc underneath and laterally."

The two phenomena described above are intimately connected with and in fact bring about another known as the "clearing of the egg". As the protoplasm is withdrawn from the center and the oil globules from the periphery, the Pipefish egg becomes "clear", i. e., the yolk, freed from these substances becomes homogeneous and translucent. At this stage the egg of Siphostoma, Fig. 1, Pl. I, consists of a button-

shaped protoplasmic disc resting on an orange-red layer of oil globules embedded in yolk and covering about one-fourth of the egg, the other three-fourths consisting of clear milky yolk. This "clearing" has been described, essentially as above, by Fusari ('90), Kowalewski ('86), and Agassiz and Whitman ('85), ^{for} Cristiceps, Corasius, and Clonolabus respectively.

In connection with the above processes, many workers, especially the students of the Salmonoids, have described amoeboid movements of the germ disc, and Vis in a recent paper ('08) has described such activities in the blastomeres up to the sixteen-celled stage. Ransom ('05) has also figured and described amoeboid movements in the yolk of Gasterosteus. These movements seem to assist in freeing the yolk of protoplasm, and the germinal disc of yolk. The opacity of the egg, which prevented my making out much about the "streaming", operated here against the detection of such movements. Once or twice, however, I thought that I did make them out, and, in one hardened germ, I found such a protuberance as is figured by Henneguy ('80) in Trout germs hardened in chromic acid.

The oil drops in the Pipefish egg are not numerous enough to make it float, but from their location they maintain the germ in an upright position. If the eggs are over-

turned, this buoyancy causes them to rotate quickly in the liquid filling the "breathing chamber" of Ponsou. How long this rotation persists I can not say, but certainly until after the closure of the blastopore. Rathke ('37) first noted this in the eggs of Black Sea forms. He also describes, as best I can make it out, an albuminous material coagulable in water or in air, which fills the "Zwischenraum" referred to above. Whatever may be the liquid filling this space in S. floridae, it does not coagulate in water, air, or in any of the fixing fluids I have used. It might be well to add here that this rotation of the egg is not a new phenomenon, having been reported, notably by Ziesler ('02) and Wis ('03) for the Salmon family.

My earliest preservations of eggs with forming germ disc were made four to five hours after the eggs had been placed in the water, hence I am not able to describe by sections its formation. In my case, however, I could not hope to add anything to the classic paper of Agassiz and Whitman on Stenolabrus, or to the more recent memoir of Behrens on the Brook Trout. Since, I preserved eggs at intervals of from five to twenty-five hours, I have sections which show the progressive degeneration of the blastodisc. So far as I know, this has never been shown and hence it may be of

interest to give a few figures illustrating this phenomenon.

Fig. 1, Pl. I, represents the sharply marked off blastodisc resting on the yolk sphere. It shows the relative diameters of blastodisc, "disque huileux", yolk sphere, and egg membrane. Fig. 28, Pl. II, is from a central section of a germ disc five hours old. The concentration of protoplasm is not yet perfect. As best I can make it out all has not yet emerged from the central yolk. The dotted line marks off a region where protoplasm and yolk are so closely intermingled as to be indistinguishable. DeLacaze ('72, Fig. 17) figures and describes a similar germ disc for the Trout. Fig. 30, Pl. II, shows a degenerating blastodisc ten hours and twenty minutes old. Such structures are not infrequent in unfertilized eggs found among others in the four- to sixteen-celled stages, in ages from eight to twelve hours. They are ^{also} found in eggs which have been in water about ten hours, and I am inclined to think, are of fairly regular occurrence in degenerating blastodiscs of unfertilized eggs.

Stricker, in 1865, describes what he called an entirely new mode of cell formation in the blastoderm of the Brook Trout, *i. e.*, a budding off of cells, which he thought originate in the amoeboid activities of the protoplasm. His figures show blastoderms with from one to twenty-three "buds".

lungs or vesicular swellings on the outer surface, and this one section is very inconclusive. Unfortunately, I have no ^{the egg of the Pipefish} showing surface views of any of these structures. The following year, Henson reported a similar cell formation in the unimpregnated eggs of the Pike. These "showed a lobulation of the concentrated formative yolk, a sort of irregular asymmetrical cleavage". After twenty-five hours, "portions of the discus proliferous were pinched off and appeared as projecting buds" This reported in 1899 that unfertilized Salmon and Trout eggs after lying in water four weeks formed hillocks on the surface of the germinal disc by the outpusing of fluid drops under the surface membrane. Neither he nor Henson gave figures. Fig. 3^o, Pl. II, makes clear these various observations.

As to the further fate of the blastodisc in the unimpregnated egg of the Pipefish, I can only say that it flattens out and finally disappears. Fig. 3^o, Pl. II, is a central section through a blastodisc twenty-six and one-half hours old, which shows this flattening. Figure 1 on the same plate shows a blastodisc taken from a lot of eggs in the invagination stage (40-48 hours). It is much larger and its lower surface is comparatively free from yolk. The contrast is evidently due to the fact that one egg has been lying free in the sea water, while the other has been under more favor-

able conditions in the aquarium. Just here it may be of interest to note that while unimpregnated eggs are often met with in the pouch with embryos of all stages, none of them ever "go bad". Ranson ('64) reports that he has kept unfertilized Trout eggs alive in running water forty-three days. More recently, His ('92) gives four weeks for the maximum ^{time} and describes the mass of germ-plasm in the unfertilized eggs of the Trout and Salmon as decreasing in size and becoming more and more set through with oil drops and yolk spheres. The degenerating diastodiscs of the Pipefish in some cases show these inclusions, but in general are quite free from them.

VI. SEGMENTATION.

Before going into a description and discussion of the segmentation of the egg of Siphostoma floridum, I wish to say that this is extraordinarily irregular. These irregularities begin as early as the two-celled stage and become very marked when eight cells are formed. The egg under consideration equals and perhaps exceeds that of the Salmon family in abnormality of cell division. The surface views were nearly all drawn from the hardened germs in 80% alcohol or xylol, the opaque egg making it impossible to draw in situ

blastodermis beyond the eight-celled stage. The drawings were all made with a Leusch and Lamb microscope and camera lucida, with the tube drawn out to 160 millimeters. The surface views were all made with the one inch eye piece and the two-thirds objective. Sections were drawn with the two inch eyepiece and the one-sixth objective.

One-Celled Stage.

This is shown in Fig. 1, Pl. I from above, and in Fig. ³², Pl. II, in section. It is high-arched, and falls steeply into the outer periblast (o.p.) from which it is clearly marked off by the circumferential furrow of the outer cell. This furrow is sometimes so pronounced in the germ disc of the Salmon family, that ^{latter} ~~the~~ λ literally overhangs its base. See Vis ('08, fig. 1) for the Trout and (fig. 2) for the Salmon. Kupffer ('06) however, says that in Syngnathus (species not given) the germ disc is not sharply marked off from the periblast, and that this condition holds till the end of the four-celled stage. Most workers on the Salmonoids, Fehrens ('06) and notably Vis ('08), represent the unsegmented blastodisc as somewhat sunken in a saucer-shaped depression. In the Pipefish, however, the blastodisc underlain with oil globules rests on a slightly flattened

area at the upper pole. Below, it is not sharply marked off from the yolk, but across its base extends a band, about as wide as the periblast to the right, composed of mixed yolk and protoplasm. The section shows several vacuoles to the right, which in the living egg were probably filled with oil. Brook ('86) describes in the Herring a blastodisc with yolk bast, Wis ('03) the like in the Salmon.

This blastodisc was found in a batch of eggs in the eight- to sixteen-celled stage (8-12 hours). Wis ('04) says the germ disc in the Salmon is formed in from one to four days. Hertwig ('03) says that the formation of the germinal disc in the Herring takes place in two hours, and in the Trout from seven to eight hours. Evidently the time varies with the kind of fish, the temperature, and the purity of the water. In the Pipefish I have found it to take place in from four to six hours. It is noteworthy that in none of the blastodiscs which were sectioned, have I ever found a nucleus. Brook ('86) could find no nuclei in the Herring until after the appearance of the third furrow.

Two-Celled Stage.

As in Teleosts generally, the blastodisc elongates slightly before the appearance of the first furrow, and, as

As a result, one axis is somewhat longer than the other. This is shown in Fig. 2, Pl. I., the normal two-celled stage, in which the blastomeres are equal. In Fig. 3, however, we have an irregular segmentation, with one cell much larger than the other and with a vacuole in the line of division. Of this type quite a number were found.

Fig. 33, Pl. II., shows a flat two-celled blastoderm not definitely marked off on the right from the outer periplast, in which the nuclei have divided. The external furrow has formed, but the cell wall has not yet come into existence. In the line of division, the protoplasmic reticulum has formed a very delicate network of dendritic fibrils arranged transversely to the plane of cleavage. Cellacher ('72, Fig. 20) describes and figures a section through two cells of a four-celled stage in the Troch front very like this. He says an indistinct streak made up of faint granulations runs vertically from the external groove towards the base. Henneguy ('80, Fig. 60) gives a figure of a two-celled stage very like Fig. 33, Pl. I., and says that the fine line dividing the two cells is bordered on each side by clear protoplasm which is traversed by very fine lines parallel to each other and perpendicular to the median line and that these fine lines lose themselves in the surrounding protoplasm. His

('98, Figs. 7, 8) illustrates and describes similar structures in the syncytium at the base of the Trout germ in early stages. In Fig. 34, we have a high arched two-celled stage in which the perfectly distinct cell wall is interrupted by a vacuole near its center. This is plainly a derivative of Fig. 24, as the preceding is of Fig. 23.

Fig. 35, is a section through Fig. 3, Pl. I., in the plane a--b, which shows the split between the two cells dilated into a large vesicle at the bottom. Very frequently the division between the two cells takes the form of a deep cleft with nearly vertical walls, and at the bottom the cleft may or may not dilate to form a small vesicle. These structures are shown in Fig. 32, and are often times much larger than figured here. In Fig. 37, we see the split being formed by the breaking down of the walls of a series of vesicles placed vertically over one another in the center of the blastoderm. This formation of vesicles in the line of cleavage was, so far as I know, first figured and described for the Trout by Oellacher in 1872. Palfour ('78, Pl. I., Figs. 6, 6a, and 6b.) figures and at some length describes vacuoles in the early furrows of the Skate. He describes such a beaded structure as shown in my Fig. 50, and thinks that they are more common than supposed, ^{and} that they play a

considerable part in the segmentation. Brook describing the
 like in the Herring but gives no figures. Kowalew-
 ski ('08, Pl. XVII, Fig. 7.) finds vesicles at the bottom of
 the furrows in the early stages of the Goldfish. Agassiz
 and Whitman ('30) figure, in surface views of Blastoderms of
Otenolabrus, rows of small vacuoles extending along the whole
 length of the cleavage planes in the two and four celled
 stages, but do not refer to them in their text. Rusari ('00,
 Figs. 4 and 5, Pl. III.) figures in both surface views and
 sections blastoderms with vacuoles. Some of the latter show
 vacuoles with large dilatations at the bottom like those in
 Figs. 30, and 31, Pl. II. § In the Pike fish, the first furrow
 does not cut through to the yolk. (See Figs. 34, 35, 36, 37).
 In this respect it agrees with Cristiceps (Rusari, '00), the
 Herring (Brook '00), Carassius (Kowalewski '08), the Bass
 (Wilson '01), the Salmon and Trout (Wis '03), but ^{is} unlike
Merluccius (Wingsley and Conn '02), Gadus (Cunningham '08),
 and others, which do cut all the way through. Agassiz and
 Whitman ('30) show that in Otenolabrus the first furrow may
 or may not penetrate to the yolk. There is never any such
 under furrow as the Bass and Otenolabrus show in the first
 division.

The eggs are laid at night, as early as ten o'clock and

probably at an hour thereafter. At any rate, by seven o'clock the next morning, they are to be found in stages of from two to sixteen cells. Probably from four to six hours since elapse before they begin to segment, it takes this long for the germ disc to form on eggs in water, in comparison with six and one-fourth hours for the Herring (Brook '02) and twelve to thirteen in the Salmon (Hoffmann'05).

Four-Cell Stage.

In Fig. 4, Pl. I. is shown a normal four-celled blastoderm. The second furrow is horizontal and crosses the first approximately at right angles. Thus there is formed a four-celled symmetrical blastoderm. Sections of this would in no wise differ from those for two-celled stages, save in the plane a--b where the beginnings of the segmentation cavity and the periblast would be found. Such a section is not at hand unfortunately.

Fig. 5, Pl. I., a more common form, shows slight inequalities in the size of its blastomeres. Such irregularities become more pronounced until they result in reniform blastoderms as Fig. 6, Pl. I. Fig. 5a, Pl. II. is a nearly horizontal section through the base of such a form as Fig. 4, Pl. I. The wide separation of two of the cells is an artefact. Of special interest are the segmentation cavity in

the center and the remnants of protoplasmic bridges which connected the blastomeres.

Eight-Cell Stage.

Into the blastoderms of the Pipefish egg of this stage many very great and seemingly irreconcilable irregularities enter and greatly confuse the investigator. These were first noted on living eggs with four and eight cells below, two, three, and four above. Hardened eggs showed the same irregularities. Surface views of a great many of these eight to sixteen-celled blastoderms were drawn. When a comparison of these drawings was made, they were found to conform to four general types. This was confirmed by an examination of all the eggs of this stage which had been preserved. At the close of this section, there is appended a table showing the relative numbers of these various types.

In Fig. 1, Pl. I., is shown the normal top of eight-celled Teleost blastoderm. It is formed by two furrows nearly parallel to the first and perpendicular to the second plane of segmentation dividing such a form as Fig. 2, Pl. I. into eight blastomeres. In this blastoderm, and in nearly all others of this and the next stage, a considerable elongation is noticeable.

Figs. 8, and 9, Pl. I., show variations of this normal type, which are more common than the type itself, but are easily referable to it. Fig. 10, Pl. II. shows a section of Fig. 7, Pl. I. in the plane a--b. In it one of the two central cells is completely cut out of the protoplasm, while, at the ^{outer} end of the cell wall, partly cutting out the other ^{cell}, there is a little split which in sections nearer the center will push a short distance to the left but on the right will extend clear across, completely cutting out and extending the segmentation cavity (s.c.). The layer of protoplasm with the yolk marked c.p. is the central periblast and the cavity above it is the segmentation cavity. This, however, is not the first appearance of either since a section in the plane a--b in Fig. 4, Pl. I., would show both. I regret that I have not been able to find such a section. The outer periblast (o.p.) never shows the periblastic ridge figured by Wilson ('91) for *Serranus*. Fig. 45, Pl. III., is through the plane a--b of Fig. 16, Pl. III., a normal sixteen-celled stage, but it will show the state of things in the plane c--d through Fig. 7, Pl. II. In this part of the normal blastoderm of this stage, the central cells are separated from the periblast (c.p.) by a large segmentation cavity, which extends for a short distance under the peripheral cells, in

this case the end cells of Pl. 7, Pl. . .

Fig. 41, Pl. II., is a section at right angles to the long axis of a blastoderm similar to Fig. 7, Pl. I. Here the two cells are separated from each other by a wide segmentation cavity (s.c.) roofed over by a protoplasmic bridge (p.b.) connecting the two blastodermis. A thin slit extends for some distance under each cell and partially separates it from the central periblast (c.p.) which is heavily laden with yolk in its lower parts. Such protoplasmic bridges as the one shown here are not uncommon in this and the next stage. All that can be said of their origin is that they have been left behind when the cells were cut out of the protoplasm. Structures similar to this would be found by making sections at right angles to Figs. 6 and 9, Pl. I. So far as I know, these protoplasmic bridges have not been figured and described before.

The periblast never comes away freely from the yolk, but is so obscured with fragments of this latter, that it has in all cases been drawn schematically, the general course of the break only being followed.


Fig. 19, Pl. I. shows a type of eight-celled blastoderm far more common in the Pipefish than the preceding. In this the plane of the third furrow shifts until it becomes equi-

torial and cuts off four somewhat smaller blastomeres from four underlying larger ones. Kennedy ('30, Fig. 30) figures a blastoderm for the trout which is almost an exact counterpart of this. A section through this blastoderm in the plane a--b reveals the structure shown in Fig. 42, Pl. II. Here the two central cells stand above the basal ones, with the line of demarcation on the left especially sharp. The segmentation cavity (s.c.) and the central periblast (c.p.) are both very much reduced.

Another very common form of eight-celled blastoderm is shown in Fig. 11, Pl. I. Here there are six cells below and two above. This is evidently a derivative of a six-celled stage frequently met with, in which two of the blastomeres of Fig. 4, Pl. I. divide by vertical furrows, the other two cells undergoing no change. Later, however, a division of these in a horizontal plane would give the structure shown in Fig. 11. Variations of this type are frequently due to the shifting of this pair of upper cells. Such a divergence is shown in Fig. 12, Pl. I., where these two cells reduced in size are shifted to one end of the longer axis of the blastoderm. Sometimes, these two cells are placed parallel to the main axis but over one of the central lateral cells. Again they may be shifted to lie at right angles to the long axis

over one^{of} the furrows separating two lateral cells so that one cell is at the edge of the blastoderm. In order not to multiply figures there is given only one drawing of sections from such blastoderms. Fig. 40, Pl. III, is a section through such a structure as Fig. 12, Pl. I, in the plane a--a. Here one central cell is very much higher than any of the other cells. The other central cell is completely cut out of the protoplasm and is roofed over by a protoplasmic bridge extending from the high cell to the left outer cell. Following the sections to one side of this, the bridge and the cell under it are found to unite. They would thus seem to have been split apart from the same mass of protoplasm.

Another eight-celled blastoderm, quite as common as either of the foregoing, is represented in Fig. 13, Pl. I. Here one cell has, by an equatorial furrow, become cut out to lie slightly above the rest. Its origin from the lower left cell is very evident. The right side of the structure is normal save that the third cell is slightly flattened at its inner edge by contact with a central cell. As in the preceding case, so here there may be variations in the position of this high level cell. It may lie in the center, at the edge, or at any intermediate position on the blastoderm. A section through the plane a--b of Fig. 13, would give a



structure essentially like that shown in Fig. 43, Pl. III., omitting the protoplasmic bridge. Mein ('72, Figs 5 and 6, Pl. XVI) shows essentially the same structures in the same stage of the Trout germ, as does Mennegy ('86) in his Fig. 35, Pl. XVII.

Fig. 14, Pl. I., is a seven-celled form, in which a unmistakable equatorial furrow has cut off three upper from four lower cells, of which three are very large. A view of this Blastoderm ^{from below} is shown in the next figure (Fig. 15). Here the two meridional furrows show quite clearly, but there is no trace of the third or equatorial furrow. The segmentation cavity, (s.c.) is so small as to be almost negligible. Unfortunately no section of this figure can be given but a comparison between it and Fig. 43, Pl. III. will make clear its internal make-up.

These nine figures of the eight-celled stage have been introduced to show; (1), the great irregularities which enter into the segmentation of the Pipefish egg at this stage; (2), that these all result from the position of the third furrow, which, ordinarily, meridional and parallel to the first and perpendicular to the second plane of division, here become equatorial; and (3), that the irregularities thus resulting may be reduced to four types which may be traced to

to the very close of segmentation. In order to illustrate definitely these points, a table is given below, and relative numbers of the different kinds of eight-celled blastomeres which have been counted.

From these eight-celled blastomeres are derived four types of segmentation which persist to the close of segmentation. From Figs. 7, 8, and 9 come two types of flat structures; from Figs. 10, 11, 15 (with the eighth cell in center), there comes a right-arc. of type of blastoderm; and from Figs. 12 and 13 -- with the eighth cell at one end -- a type of blastoderm thick at one end and tapering toward the other. These structures will be more clearly shown in the next section.

TABLE TWO -- RELATIVE NUMBERS OF BLASTOMERES OF EACH TYPE OF EIGHT-CELLED STAGE OF THE PIPPIA EGG.

Figures refer to the figures on Pl. I.

Kind in	Fig.	II, VIII, IV.	V.	VI.	VIII.	IX.
Paronyi	1	2	2	3	2	0
Formalin	2	2	3	4	4	0
Paronyi	3	0	3	1	4	1
Formalin	4	7	7	3	3	0
Formalin	5	2*	2	3	3	0 + 6-celled.
Sub-acetic	6	0	0	2	4	0
Paronyi	7	2	1	0	2	0
Total	7	8	15	20	22	0

Sixteen-Cell Stage.

Intermediate between the eight and sixteen-celled stages, are found many blastoderms with twelve, fourteen, and fifteen cells. These are in fact more abundant than blastoderms with exactly sixteen cells.

Figs. 15 and 17, Pl. I, show the two most regular sixteen-celled stages that have been found, yet they do not have the regular structure of the corresponding stages shown for Gerrhus by Wilson ('11) and for Cristiceps by Fusari ('30). These blastoderms have been formed by each of the cells in Figs. 7, 9, or 10, Pl. I, dividing into two. In Fig. 15 all the cells save one are practically on the same level, or at most with a gentle curve across the upper surface. In Fig. 17, the blastoderm is arranged more irregularly. Fig. 18, Pl. II., is a section in the plane a-b of a blastoderm like Fig. 15, Pl. I., preparing to divide into thirty-two cells. The two central cells will divide to form two surface and two interior cells, while the outer cells will each divide into two cells both on the surface. This is shown by the position of the centrosomes. The cells form a gentle arch roofing over a considerable segmentation cavity. The planes of segmentation are dilated at their outer ends into vesicles which are covered by thin protoplasmic sheets or bridges. Fig. 19,

Pl. III. is a section of some such structure as Fig. 17, Pl. I. in the plane a-b. Some blastoderms of this stage have been found in which the four or five cells were not cut off from the basal periblast, but these are too infrequent and too little understood to be reproduced here. Fusari ('30) has figured a section like this for Crisidopsis, a Gony.

In Fig. 46, Pl. III., there is shown a section of a flat-topped, abrupt-edged, sixteen-celled blastoderm of a type which persists till the preparation for invagination begins. What the appearance of such a blastoderm in surface view would be, I can not say; probably it would in no wise differ from Fig. 13, Pl. I. The essential difference between Figs. 44 and 46, Pl. III. is the circular groove sharply marking off the outer periblast (o.p.) in the latter. Possibly these figures are derivatives of the one-celled stages shown in Figs. 10 and 13, Pl. I. In the figure in question, there is a large segmentation cavity (s.c.), and a yolk-laden periblast (o.p.). The dotted lines show where the outer periblast has been torn away. Note the large dilatation at the outer end of the right furrow and the protoplasmic bridge covering it. Fig. 45, Pl. I., is a derivative of some such forms as Figs. 10, 11, 13, and 14 of the same plate. It is arched but the crest of the arch is not in the center but to one side and the cells lie

in two if not three levels. A section through an almost identical form in the same plane q-r, is shown in Fig. 12, Pl. III, and makes clear its sloping outline, and its two eccentrically placed high cells. It has one interior cell, which in the next section is clear of the central periblast (c.p.) and has probably originated by the horizontal division of an outer cell.

Fig. 13, Pl. I., shows a modification of the arched type. Its sixteen cells are in two layers, and the seven upper cells are on an approximate level. Fig. 14, Pl. III, is a section through some such blastoderm as this one. Its surface slopes gently, and the left peripheral cell projects over the outer periblast (o.p.). This latter ^{structure} will be found frequently in later stages. Vaguettes are found in two of the division walls.

The high-arched type of sixteen-celled blastoderm is shown in Fig. 20, Pl. I. This is probably a descendant of a blastoderm like Fig. 14 on the same plate. No description of it is needed, beyond calling attention to the fact that the five upper cells are cut out by an equatorial furrow. This is seen by referring to Fig. 21, which is a ventral view of the same blastoderm. Here only five of the vertical planes seen from above cut all the way through. The

one marked g in Pl. I, has not reached the base. The small segmentation cavity (s.c.) recalls that of Pl. 14. Let us compare with this figure the next, No. 22, which is a view from below of a similar high-piled sixteen-cell stage. Here there are nine basal cells resting on the yolk, six in the second tier, and a central one forming the keystone of the arch, the whole enclosing a spacious segmentation cavity. Bearing the fact that the segmentation cavity (s.c.) extends under the marginal cells, Fig. 4, Pl. III, may be given as a section through Pl. 22 in any plane passing through the keystone cell. The central cell has not yet completely cut itself off from its neighbor to the right, and the cell to the left has a resting nucleus obviously elongate.

There have now been figured and described in surface views and sections, such sixteen-cell structures as may be considered typical for the Pipefish. Of these two are sufficiently like the usual talcost forms as to be called normal, but a great majority, fully ninety percent of those studied, are like figures 13, 13, and 23, Pl. I. In this connection, Hertwig's statement (Handbuch, pp. 543-4) with reference to the fourth segmentation and formation of the sixteen-celled stage, is of interest. He says, "The end result is everywhere the same, a 'check-board-like' arrangement of sixteen

blastomeres, four in the center, and a circle of twelve equatorial cells". To construct this is for the Pipera, proceed at the figures given, and at the table shown on page 55 will demonstrate.

Equatorial Plane of Segmentation.

All investigators are agreed as to the homology between the first and second furrows in teleost^{an} and amphibian eggs, but whether or not the third furrow corresponds is a very debated question.

Hoffmann ('31) figured and described in pelagic fishes the first segmentation as equatorial, dividing germ from periblast; but, later ('35) he acknowledges his error and declares that in Salmo the third furrow is equatorial. Ziegler says that the third furrow in the Salmon and trout is equatorial and divides eight upper from eight lower cells, the latter not being as yet marked off from a periblast. Rauber ('83) made a careful study of the subject based on the well known fact that the fourth amphibian furrow in a great many cases is not truly meridional but avoids the pole and forms many structures like Figs. 5 and 6, Pl. I. He concludes that the first equatorial furrow of the frog has been lost in the Teleost, and homologizes the third teleostean

e

f

g

h

①

②

③

④

⑤

⑥

⑦

⑧

⑨

⑩

⑪

⑫

⑬

⑭

⑮

⑯

⑰

⑱

⑲

⑳

㉑

㉒

㉓

㉔

㉕

㉖

㉗

㉘

㉙

㉚

㉛

㉜

㉝

㉞

㉟

㊱

㊲

㊳

㊴

㊵

㊶

㊷

㊸

㊹

㊺

㊻

㊼

㊽

㊾

㊿

Ⓚ

Ⓛ

Ⓜ

Ⓝ

Ⓞ

Ⓟ

Ⓠ

Ⓡ

Ⓢ

Ⓣ

Ⓤ

Ⓥ

Ⓦ

Ⓧ

Ⓨ

Ⓩ

ⓐ

ⓑ

ⓒ

ⓓ

ⓔ

ⓕ

ⓖ

ⓗ

ⓘ

ⓙ

ⓚ

ⓛ

ⓜ

ⓝ

ⓞ

ⓟ

ⓠ

ⓡ

ⓢ

ⓣ

ⓤ

ⓥ

ⓦ

ⓧ

ⓨ

ⓩ

⓪

⓫

⓬

⓭

⓮

⓯

⓰

⓱

⓲

⓳

⓴

⓵

⓶

⓷

⓸

⓹

⓺

⓻

⓼

⓽

⓾

⓿

Ⓚ

Ⓛ

Ⓜ

Ⓝ

Ⓞ

Ⓟ

Ⓠ

Ⓡ

Ⓢ

Ⓣ

Ⓤ

Ⓥ

Ⓦ

Ⓧ

Ⓨ

Ⓩ

ⓐ

ⓑ

ⓒ

ⓓ

ⓔ

ⓕ

ⓖ

ⓗ

ⓘ

ⓙ

ⓚ

ⓛ

ⓜ

ⓝ

ⓞ

ⓟ

ⓠ

ⓡ

ⓢ

ⓣ


ⓤ

with the fourth (pole-avoiding meridional) furrow of the frog. For this interpretation of Hamber see Wilson ('91, pp. 214-16).

Agassiz and Whitman ('90) think that the amphibian equatorial furrow has become vertical in the teleost, and that the horizontal division of the four central cells of the sixteen-celled stage into four outer and four inner lying cells is the first equatorial segmentation. With this latter statement Topser ('01), from his work on *Salmo*, is in full accord. Brook ('06) describes, from sections of *Ferrugia* eels (Pl. XIII., Fig. 9), an equatorial segmentation separating the four blastomeres from the periclast. List ('07, Pl. XXVI., Figs. 4 and 5) finds the second furrow in *Grenilabus* to be equatorial, and says that Kupffer found the same in the *Ferrug.* In *Cristiceps*, Fusari ('30, Figs. 4 and 5, Pls. I. and III.) finds that in the sixteen-celled stage, all the cells are united at the base, but the next sets ^{sets} division, sixteen central cells free from the yolk and from sixteen peripheral cells. This he calls the equatorial division. Wilson ('91, p. 215) agrees with Hamber in all points. Salsassa ('36), in the segmentation of *Callionectids*, finds as a rule that an equatorial division follows the eight-celled stage, although it sometimes comes earlier.



An equatorial segmentation has been pointed out in certain eight-celled blastoderms of Siphonura, and this gives them a very decided resemblance to the upper surface of dividing amphibian eggs. Grönroos ('00) (See Herwig's Handbuch) gives a series of figures for the eggs of Tritons, to which, the figures above note, show very striking resemblances. The Tritons have eggs with relatively large amounts of yolk, and in them the segmentation approaches the meroblastic condition. The text-figure reproduces some of the more striking forms to which reference will be made. The resemblance is so striking that no extended comparison is called for. With Grönroos' Fig. I., compare Fig. 17, Pl. I, and also Meneguy's ('00) Fig. 33. They are almost identical. For a figure which almost duplicates his Figs. II, and III, see Fig. 18 on the same plate. Another drawing not included in the plates, is one almost identical with his Fig. IV, and Figs. V, and VI, are very similar to Fig. 3. The comparison might be extended further, but this is sufficient to show the very striking similarity between these two forms. That we have here an analogous segmentation is beyond question. The segmentation in the Pipefish egg in the blastoderms in question, is equatorial, or at least approaches very closely thereto, and it seems hardly going too far to say that such



Pipefish *Blastodermis* as Figs. 13, 14, 15, 16, 20, Pl. I., there is a reversion to a type of segmentation essentially like that of Amphibia.

Thirty-two-celled Stage.

Normal types of this stage, as shown in Fig. 23, Pl. I., were found to make up about thirty percent of one lot of eggs, and were noted sparingly in all others. Fig. 23 is plainly a derivative of forms like Figs 16 and 17, and, while it may be called normal, is noticeably different from Wilson's figures of the same stage for *Serranus* (Pls. 8, 9, 9, and 10). No section of this stage will be given. Its internal structure will be made clear by reference to Fig. 42, Pl. II., a sixteen-celled blastoderm ready to divide into thirty-two. The two central cells will divide horizontally, the two lateral ones by an oblique plane resulting in six surface and two interior cells. (Compare Wilson's Fig. 10)

Fig. 43, Pl. III, is a section from a flat-topped acropod-edged blastoderm, drawn with the same magnification as the others. It serves to show the inequalities in the size of the blastodermis. The peripheral cells are very much flattened above, though retaining their rounded forms below. To the right the section cuts the point of a sixth cell. The

segmentation cavity (s.c.) is partially filled with cells. The larger and lower ^{cell} seems to have been cut off from the central periblast (c.p.), from which it is separated by a cell wall so delicate that the oil immersion only will detect it. It is like the periblast further in that its periphery contains many yolk granules.

Fig. 24, Pl. I., is an arched type, with the highest point rather nearer the lower side. The twenty-seven outer cells are in three tiers, and while the second is pretty sharply marked off from the first there is but little difference in level between it and the third tier. There is here noticeable a symmetry comparable to that in Figs. 7 and 8. The plane 1-1 in all probability represents the first, 2-2 the second line of division referable to Fig. 4.

A central section through Fig. 24 in the plane a-b is shown in fig. 25, Pl. III. The peripheral cells form an arch with the highest point slightly to one side, and enclose a segmentation cavity which is almost filled with cells. The two smaller cells have been cut off from the peripheral ones, the larger probably from one of its fellows. The periblast (c.p.) is thick and yolk. A more pronounced large-ended type is Fig. 21. Here the segmentation cavity is somewhat eccentric, and, as in the preceding, the thick end over-

names the base. The spacious segmentation cavity (s.c.) contains one cell which warts on a curious tongue of protoplasm from a partially segmented region on the left.

Fig. 52, Pl. I., is a typical nine-celled blastoderm, whose cells are arranged in three layers. Its highest cell x is slightly eccentrically placed, and one of the axes is somewhat longer than the other. Fig. 52, Pl. III., is a central section through a similar but slightly older blastoderm. The marginal cells are sharply marked off from the outer periblast (p.r.). The arc is high and round. On the left three cells are imperfectly separated, and from one of them a tongue of protoplasm, from which a cell has been cut off, projects into the large segmentation cavity. The periblast, torn off at the right, is in the center reduced to a mere film of protoplasm with mucopol. adherent below, thus giving it the breadth as drawn.

Fig. 53 shows a structure by no means uncommon in the egg of Ciphostoma. It is a thirty-two-celled stage in which no periblast has yet been formed. The cells are in two layers, the long cell on the upper right is nearly ready to divide, and underneath the whole is a thick layer of protoplasm in which three vertical cell walls extend downward and are lost. Later transverse walls will appear and cut these

cells out of the spectrum, finally leaving a peripheral layer below. There is a very small segmental cavity (s.c.) and the large cell to the right has a vacuole (v.). Fiedler ('52, Fig. 2) figures an almost identical structure for the Salmon. Nowalewski ('56, Figs. 1 and 2) portrays essentially the same conditions in the Goldfish. Hoffmann ('55, Figs. 6 and 9 especially) describes a similar structure in the Salmon germ. And latest of all, '53 ('53, Figs. 7 and 10) confirms the figures and descriptions of the earlier workers on the Salmonids.

Fig. 23, Pl. I., is a very interesting divided blastoderm of this stage with sixteen cells in one division and fourteen in the other. Such structures have been met with occasionally in stages of from sixteen to sixty-four cells, but especially abundant in the eggs from one fish. Out of twenty of these eggs killed in micro-acetic, five were like the one figured. That these were not artefacts is shown by the fact that eggs of the same lot killed in formalin also contained divided blastoderms, the numbers of which were unfortunately not noted. In each division a segmental cavity exists and the line of separation is broad and definite down to the periblast. These points are brought out very definitely in Fig. 54, Pl. III., a section through a similar but older blasto-

form. In the left half there is a small separation cavity (s.c.), on the right, however, there is none.

There is no periblast. Cells have been cut out of the mass of protoplasm, leaving a thick germ basis in which are found vertical cell walls and a number of vacuoles (y.), and which is filled below with fragments of yolk. Fig. 39 Pl. IV., is a divided sixty-four-celled stage of the thick-ended type. The distance between the two parts is here not so wide. In other cases they swell out to a vesicle at the bottom or are reduced to a mere line as in the two-celled stages above. There is a segmentation cavity in each portion, but there is no distinct periblast, the basal layer of protoplasm being thick with a large vacuole and full of yolk in its lower part. In some cases where the plane of separation is reduced to a line, the cells are drawn out into long points toward the base as if a fine thread, used to separate the parts, had elongated the cells upward.

The only reference to such peculiar conditions as shown found in these figures is in a short section on *Corythoeca* in Eycleshymer's paper on *Amphistoma* ('35, Fig. 30, and others). This writer thinks, however, that these divided blastomeres do not result in double embryos. The same seems to hold



the true for Pipefishes of Beaufort, for although thousands of eggs and larvae and hundreds of adults, alive or preserve, have been examined, only two apparent cases of deformation have been found by the writer. The literature of these fishes contains no few references to abnormalities. H. Stein ('62) described a *Syngnathus* with two caudal fins. ('64) reports a *Syngnathus* with two heads. However, Gault ('67) reports in *Syngnathidae* of the Black Sea many abnormalities of the snout, eyes, and tail, but he thinks the retardation or development.

A fair example of the late stages of segmentation is shown in Fig. 27, Pl. I. Here the least-eight cells are in three tiers, with one cell high above all. There is a slight deviation in one axis, possibly a derivation of the condition found in the eight-cell stage, and a curious, regular arrangement of certain cells. On the whole, however, the segmentation is very irregular, and it becomes more so later, finally all trace of symmetry is lost, and the blastoderms become almost circular in outline. No surface views of later stages will be given, since, as the cells grow smaller, the blastoderms approach more and more the ordinary teleostean form.

Stage of Sixty-four Cells.

Artificial fertilization being impossible in *Siphonocera*, one can not divide late meroblast into stages by hours, and the presently varying shapes of the blastoderms make it impracticable to classify sections by the number of rows of cells in each, as some writers do, so it has become necessary to devise an arbitrary scheme. This scheme is to count the peripheral cells in the central section of a blastocyst, then, assuming a like number in a section at right angles to this, by squaring this number, the approximate number of surface cells is found. The size of the cells serves as a check to this.

Fig. 36, Pl. IV., with eight peripheral cells, is from a normal type of λ sixty-four-celled stage. The periblast (p.b.) is thick and yolk, and at the right is a cell not yet cut off from it. The segmentation cavity (s.c.) is filled with cells, some of which are ready to divide.

Fig. 37 is ^{derived} from a flat blastocyst of the λ stage, and, ^{preceding} in comparison with Figs. 43 and 46, Pl. II., is seen to have undergone considerable division, in horizontal planes, as is shown by the number of cells filling the segmentation cavity. The large nuclei ^{are} in the spirose stage, and in the left marginal cell there are two large vacuoles.

The high-arched type of this stage is shown in Fig. 38,

a derivative of a structure like Fig. 33, 34, III. The surface falls steeply into the outer periblast (o.p.), the cells are all rounded and have small nuclei. Very interesting are the two cells which are inconspicuously cut off from the central periblast (c.p.). Scattered yolk granules are found in some of the cells. The mitotic figures indicate that division into the next stage has begun.

In Fig. 33, we have an example of the thick-ended type. The section is slightly to one side of the center, and shows one cell just free and another not yet cut out from the thick yolk periblast. Note the vacuoles which help to delimit cells. In the central section the small segmentation cavity (s.c.) becomes somewhat larger. The outer cells are flattened on the exterior, and the whole structure is very like Fig. 32.

Stage of One-hundred-twenty-eight Surface Cells.

The normal gently arched type is represented in Fig. 30, a nearly central section of a blastoderm of this stage. The central periblast (c.p.) is here thick and fairly well delimited from the yolk below. Of especial interest are the cells in the act of being cut out of it into the segmentation cavity. Very notable is the agency of vacuoles (v.) in this

process. The cell next to the right marginal cell was in its later part a nucleus, the first cell within the second st region.

Fig. 31 is an example of the first-arched type. The central peripheral cells, like those of the preceding stage, have undergone more division than their fellows. The periblast at the left is reduced to a mere line, at the point it is thicker and so filled with yolk that one can find no line of separation save where the whole has come away from the yolk.

The round-arched type finds a good illustration in Fig. 32. There are three points of interest in this section: the presence of vacuoles which help to separate the right marginal cell from the "Blast"; the cell near the center still adherent to the central periblast, and, with its neighbors, having some yolk particles in it; and two pairs of neighboring cells with spindles at right angles to each other. These last illustrate the exceedingly irregular segmentation in the Pipefish egg.

Fig. 33 is a nearly central section through a blastoderm intermediate between the normal and the thick-ended type. It is sharply marked off from the outer periblast which it overhangs on the right. The segmentation cavity is reduced to

the interstices between the cells. All along the germ axis in all the sections cells are being cut out and the peripheral layer left behind. An especially interesting case of this is found in the very center. Some cells show mitotic figures, and in others there are beside the nuclei small solidly staining round bodies, of unknown function.

Fig. 34 derived from Fig. 33 is a fine example of its type. It is very flat and the segmentation cavity is very much recessed. The periblast, perfectly free from yolk and as distinct below as above, has a layer of cells cut out of it and at the left a nucleus under the marginal cell and clearly derived from it. At one point near the center, the periblast is recessed to a mere line. This figure, which is typical for the whole blastoderm, is remarkably like Wis' ('38, Fig. 17) for the Brook Trout.

Stage of Two-hundred-fifty-six Surface Cells.

The normal type blastoderm of this stage is shown in Fig. 35. The cells lying near the upper surface are considerably smaller than those in the lower parts nearer the periblast. To right and left are furrows with dilations helping to cut cells out of the periblast, and at the center are cells nearly free from it.

Fig. 66 is plainly a derivative of Fig. 63, in its general outline and in the peripheral series which separate its outer periblast (o.p.) from the marginal cells. The periblast is somewhat sunken in the yolk and free from cells throughout the whole blastoderm. The segmentation cavity, because of this depression, large and is only partly filled with cells. Neighboring sections show the upper surface to be as flat as that in Fig. 61.

The third type is shown in Fig. 67 from a nearly central section. There is a very noticeable difference in the size of the blastomeres, some being fully three times as large as others. Here again are cells being cut out of the basal periblast. They are in all stages from rounded buds to completely cut out cells. Neighboring sections show nuclei in each of these. At the right are two cells connected by a stout protoplasmic bridge.

Fig. 64 Pl. 7., is a good example of the rounded type. The spacious segmentation cavity is loosely filled with rounded cells. The periblast is throughout the blastoderm in the form of two thick pads in the border region, but in the center it is very thin and obscure with yolk. Nowhere in the whole blastoderm are cells being budded off from it. In the peripheral cells,

there are even in this advanced stage, but gaps of protoplasmic bridges.

A nearly horizontal section through such a blastoderm, as Fig. 60 is shown in Pl. IV. This is introduced to show the arrangement of cells in horizontal plane. There is here a closer aggregation of cells to the periphery, the inner row being a derivative of the outer, while in the center the cells are more scattering.

Fig. 60 is from a blastoderm intermediate between those from which Figs. 58 and 57, Pl. IV., are taken. Neighboring sections are more like Fig. 60. Some of the outer cells show a tendency to elongate and are somewhat smaller than the interior ones. Both marginal pads are nucleated, and in one a cell wall is cutting downward. While the periblast has cells resting on it and even depressing it, nowhere in the blastoderm is there any evidence that one have been budded off.

Stage with Five-hundred-twelve Cells on the Surface.

Fig. 71, the normal type, is very similar to the preceding figure. Here the cells are pretty uniform in size, and those on the surface are noticeably elongated, some being drawn out in fine thread-like connections — the begin-

ning of the "Hörschicht" of von Cuvier. Some of the nuclei are in process of division by mitosis, but the majority stain solidly. The outer thickening of the periblasts are nucleate, the basal portion is thin, pale, and usually devoid of either nuclei or cells.

The rounded type is finely shown in Pl. 92. The surface cells are slightly flattened and only occasionally pointed, and one on the right is binucleate. The blastomeres are by no means uniform in size, and on the right is a giant cell with a proportionate nucleus. All the nuclei stain solidly. The periblast is very thick and, while laden with yolk fragments, is fairly distinct below. There are two nuclei in the periblast. One is in a thickening out of which a cell will probably be formed. Several are cells which seem to have been recently cut out.

Fig. 93 is an excellent illustration of the flat type. The blastomeres are very uniform in size and distribution, and are especially noteworthy for the large number of dividing nuclei with spindles at all angles. The chief interest, however, centers in the periblast, which is thick and possesses many yolk granules, but is perfectly distinct. In it to the right is a nucleus dividing by mitosis with a spindle considerably longer than those in the blastomeres.

On the left, the section cuts through a chromatin mass at right angles to the spindle. At the extreme left is found, for the first time, a nucleus in the outer periblast. The periblast in this blastoderm is very rich in nuclei dividing by mitosis. A cursory examination shows one vertical and eight horizontal ones. Another blastoderm, of the same lot and stage, contains, in its periblast, thirty-three oblique spindles at all angles from nearly vertical to nearly horizontal, twenty-nine lying horizontally, and seven standing in a vertical position; in all sixty-nine spindles were counted (none twice). There are a very few solidly staining nuclei, but a great number are cut, as above, through the chromatin masses, and these are not counted. There can be no doubt that the spindles stand in all positions.

The last type of this stage is Fig. 7. The cells are not uniform in size, and many are twice as large as the small ones. Most of the nuclei stain solidly, but some contain spindles. Two bi-nucleate cells are present, the one in the periphery being very large. This condition is far from rare in this and later stages. Some thirty cases have been particularly noted. The periblast is very thick, yolkly, and distinct. It contains several qu-

clei, and a cell is either being cut out or is in process of uniting with the periblast. In other sections similar conditions are found. The reentrant angle, between the outer periblast and the "band" in this and Fig 77, recalls the line in Figs. 55 and 60, Pl. IV., and Fig. 47, Pl. II, and in 'his' figures for the Salmonoids referred to above.

Stage of One Thousand-twenty-four Surface Cells.

Fig 76 represents the normal type and presents several points of interest. The surface cells show a considerable flattening and adjacent to the are other cells with their bases generally at right angles to the former, making the outer layer in places two cells thick. The inner cells show a tendency to run together in three's and four's. The chief interest however centers in the periblast. This is notably free from yolk and is drawn exactly as it appears. Nuclei are scattered very freely throughout its entire extent in all sections, and nearly surround the large vacuole to the right of the center. At the left a large cell, which has recently been cut out of the "band", is dividing by mitosis. A large number of cells rest on and indent the periblast and are either being cut out of or added to this layer. The close juxtaposition of these cells to nuclei in

the periblast would seem to lead to the former conclusion.

The second type is represented in Fig. 76, which, judging by the number of cells in the periphery and by the size, is from a blastoderm slightly younger than the preceding. The periblast is sunken deeply into the yolk and has, thus nearly doubled the segmentation cavity, which is sparsely filled with scattered cells. The thick periblast is so obscured with yolk that no nuclei could be found. It is here free from cells, but nearby sections show a condition in this respect like the preceding figure. In the Deckschicht, near the center, is a binucleate cell, while its neighbor has a spindle.

Fig. 77 is from a rounded blastoderm of about the same stage as the preceding. A Deckschicht can hardly be spoken of here, for the outer cells are nearly all round. The segmentation cavity is reduced to the small interstices between the cells. The greatly thickened periblast is full of large vacuoles and abounds in nuclei in all the sections and near the center seems to be budding off cells. In the left Rand there is a mitotic figure fully twice as large as any in the blastomeres.

No better illustration, of the lens-shaped blastoderm, so characteristic of late teleost segmentation, than Fig. 76,

can be given. It probably has been derived from a form like the preceding by the pressure of the cells against the periblast causing the periblast to be depressed. Thus the segmentation cavity has been enlarged and the cells are more scattered than in the preceding. The cells are grouped in two's, three's and four's.. The thick periblast has several nuclei in the resting condition. There is a well defined "epidermic stratum", as the English writers term the outer layer of cells.

Fig. 73 represents the last type of this stage, we need detain us but for a few moments. Its outer cells are flattened and unequal in size, and the interior cells are the largest of all. The periblast is very thick, yolk, and indented from below by large vacuoles. On the left a large cell has been cut out of the periblast and at the right a cell indents the periblast, while in the center cells seem to be in process of formation from the basal layer. This blastoderm is closely related to that illustrated in section by Fig. 74.

Fig. 80 is a horizontal section through some such blastoderm as that shown in section in Fig. 73, Pl. V. It shows the loose arrangement of the interior cells, and the drawn-out cells of the Deckschicht. This^{all} was broken at several

points, in the process of sectioning.

Latest Stages of Segmentation.

From this time on, it is not profitable and is hardly possible to follow the segmentation, but some figures may be introduced to show the course of development.

Fig. 61, Pl. VI., is probably a descendant of a form like Figs. 71 and 72, Pl. IV. There is an "epidemic stratum", the cells are loosely scattered in the large segmentation cavity. The periblast is quite distinct, free from yolk, and has a good many nuclei. Just across the border from one of these nuclei is a cell, in another place a cell lies in a depression in the periblast.

Fig. 62 is another type with ^{with} Deckenschicht, cells fairly ^{with} closely crowded in one segmentation cavity, and a very thin periblast out of which cells are being ^{with} moulded or into which they lose themselves. At one or two places, the periblast is reduced to the thickness of a cell wall, and in neighboring sections nuclei abound in it. In the right outer periblast two tripolar spindles are found. These have been noticed occasionally in other sections.

Fig. 63 is the typical teleost late lens-shaped blastoderm. It closely resembles Fusari's ('39) Fig. 3 for

Bristiceps, and is almost a duplicate of Salpasa's (1901) Fig. 3 for the Salmon in corresponding stages. The invagination of the blastoderm into the yolk is probably due to pressure against the egg shell. In the highest part of the epidermic stratum is a very large cell, and in the right hand a giant nucleus, which is separated from the neighboring cells by hardly more than the cell wall. As a cell on the left, has been cut out one of the hand. The thin periblast was resting on it many cells, neither the origin nor the fate of which can safely be guessed upon.

Early Stages Preparatory to Invagination.

Figure is a normal type in which the cells were beginning to move away from the periblast, to crowd together in the upper part of the blastoderm, and to leave a sub-terminal cavity, (s.t.c.) between them and the periblast. The line marked x is, in this and the following sections, the lower limit of the cells. The out most cells of the blastoderm have flattened until they make a very thin skin-like layer. The periblast is comparatively free from yolk granules and is here drawn after nature instead of semi-diagrammatically.

The second type is represented in Fig. 2a. The cells

are closely crowded, the periblast, composed of the sub-epithelial cells (s.e.c.) is very large. The periblast is very thin and well-laminated showing strands that only one nucleus could squeeze out.

Fig. 30 illustrates the third-order type. In this specimen the cells are not so closely crowded as in the previous, and a distinct sub-epithelial cavity is formed. The very distinct periblast consists of the s.e.c. and, on the left is separated from the blastoderm by a sharp constriction. A very large sub-epithelial cell is shown and nearly two others are shown. On the left is shown a cell of ordinary size.

Fig. 31 represents the first-order of type line 31.35, which has begun to flatten and is preparation for the next stage. This flattening is probably responsible for the small sub-epithelial cavity. The periblast has now large nuclei. Two distemperes shown indicate the size of the cells at this stage.

Late Stages Preparatory to Invagination.

Of these two, two will be shown. Fig. 32 is the normal latest structure for this stage. The cells are all

crossly drawn since a right-angle line, making a large
 sub-axial cavity (p.d.c.) below. The particles in here
 filled with gold and contains very fine iron particles. The
 dimensions are mostly in area but over the gold, in the
 section in Fig. 30 is twenty-five percent larger than in
 in Fig. 31.

When the effects of the above is given by Fig. 32 in
 comparison with Fig. 30 it can be distinguished below of the
 middle part, or rather it is the same as the middle part
 like Fig. 30 would be very obvious. Further, the direc-
 tion iron is shown. The particles is filled with iron
 fragments and the metal are very much fractured.

The Periblast.

The origin of this layer, together with many of its peculiarities of structure, has been noted in the descriptions of the plates. It is not my intention to get into any extended discussion of its formation and fate. However it will be well to describe briefly the various modes of its formation in other Teleosts and to show under which of these classes the Pipefish falls, and finally to give references to a few of the more valuable papers on this subject.

In Teleosts, the periblast layer seems to be formed after three types:

I. In eggs, in which the first furrow cuts through to the yolk, the periblast is formed by a thin protoplasmic sheet extending inward from the "Blast", Henneguy ('83, Fig. 63) shows this very plainly for the trout.

II. In eggs in which there is no layer of oil drops under the germ-disc, or those in which the protoplasmic mass separates sharply from the yolk, the periblast is formed when the inner ends of the cells in the four and eight-celled stages are cut out and lifted from the under-lying thin protoplasmic sheet. This is the mode of formation in Serranus (Wilson '81), Leptocephalus (Agassiz and Willard '80),



and Belone (Tarboen '01).

III. In eggs in which there is an imperfect separation of germ disc and yolk, or in which there is a layer of oil drops under the blastodisc, the resulting periblast has a very peculiar mode of origin. Cells are cut out of the protoplasmic disc in successive layers from above downward, and the central periblast is the remnant of blastodisc left when this process has ended. The explanation for this is that the protoplasm continues to flow out of the yolk into the germ disc until segmentation has progressed some distance. Wu ('05) noticed that the germ disc was not fully formed in Symphyla until after the four-celled stage.

This formation for the central periblast is described by most workers on the Salmonoids, notably by Wu ('05) and Hoffmann ('06) for the Salmon, and later of all by Wu ('08) for the Salmon and Trout. Kowalewski ('08) found essentially the same formation in Carassius and Polyacanthus.

The central periblast nuclei, in types I. and II., originate by division of the "Polar" nuclei and migrate centralward in this paper. In type III., they are the direct descendants of the segmentation cells.

In Siphostoma floridus, there are found the two meth-



has of central periblast formation described in types II, and III, above. In Figs. 40, 46, 47, 48, 52 for the eight and sixteen-celled stages, there is shown a mode of formation for the periblast which negatives the idea that from it there could ever come an after-segmentation. On the other hand in Figs. 53, 54, 55, 56, 59, 61, 62, the central periblast is the protoplasmic remnant of the primary germ disc, left after all the blastoderm cells have been cut out of it. It is hard to note here, that a migration of nuclei into the marginal region and the formation of a "wreath" by the disappearance of cell walls, has, because of the opacity of the egg, not been seen in the Pipefish. Whether it takes place or not, I can not say.

The difficult question, whether, in the egg of the Pipefish, cells are budded off from the central periblast and added to the blastodermes, can not here be taken up. However this would seem to be a legitimate consequence of such a mode of cell formation as that shown in type III, above, and apparently finds confirmation in Figs. 76, 77, 79, 80, in which a perfectly definite periblast layer has been formed. If these figures are compared with His' ('88) Figs. 10 and 12, this matter will be made clearer.

For a fuller discussion of the origin of the periblast and its nuclei, and of the fate of the latter, the reader is referred to Frank ('33), Howland ('33), Hoffmann ('33), Masai ('30), Sargent ('33), Zeidler ('37 and '38), His ('32), and Hartwig ('33).

At this point, the work on the development of the Pipefish will have to rest. It has been the intention of the writer to carry it further, at least to the point of the blastopore, and for this purpose the sections have been cut, and the difficulties met with have caused so many delays, and the work for the present is brought to a close here.

The eggs of the Pipefish is very different from most other teleostegans in the form of its segmentation and the actual origin of its periblast together with the "after-segmentation" of cells therefrom. So marked are these differences that it seems proper to state that the figures in this paper are representative of the sections of a thousand or more eggs, obtained from thirty-three fishes.

BIBLIOGRAPHY.

1880. Agassiz, A. and Thomsen, C. J. On the Development of Some Polar Fishes. Preliminary Notice. Proceedings American Acad. Arts and Sciences, Vol. XV.
1885. _____, _____. The Development of Osseous Fishes: II. The Pre-Embryonic Stages of Development. Memoirs of the Museum of Comparative Anatomy, Vol. XIV.
1885. Barbour, T. S. A Monograph on the Development of Elasmobranch Fishes. London.
1886. Behrens, C. Die Reifung und Befruchtung des Torfeneids. Anat. Hefte, Abt. II.
1886. Böeck, Nikolaj. Zur Kenntnis des Fortschrittes und der Keimblätterdifferenzierung in Ei der Knochenfische. Jenaische Zeitschrift für Naturwiss., Bd. XV.
1887. Brook, George. Formation of the Germinal Layers in Teleostei. Trans. Royal Society of Edinburgh, Vol. XXVI., Part I. Session 1886-87.
1890. Born, Ludwi. Wenn die Fruchtasche von Synbranchus typic. Anatomischer Anzeiger, Vol. XVII.
1897. Cunningham, J. E. On the Histology of the Ovary and of the Ovarian Ova in Certain Marine Fishes. Quart. Jour. Mic. Science, Vol. XI.

1850. Day, Francis. On the Fishes of Cochin-China, etc.
Proceedings of the Zoological Society of London.
1876. _____. The Fishes of India, etc. London.
1880. Bonn, Basileus. The Early Development of Bar-Pike
and Sturgeon. Journal of Morphology, Vol. VI.
1848. Poyé, G. P. M. Sur un Meropse dans les Oeufs
au temps media et conjugatus opinion. Société
Philomatique, ---- Extraits des Procès - Verbaux des
Séances.
1834. Schneider, Georg. Die Fische der Nordsee an Poldinsee.
Mittheilung von der Naturhistorischen in Paderborn. XVI. Jähr.
1831. Eckstrom, G. M. Fiskarne i Örnö Skärgård. Om en a
Svenska Vetenskaps - Akademiens. Handlingar. Stock-
holm.
1890. Huxley, A. C. The Early Development of Ambly-
stoma with Observations on Some Other Vertebrates.
Journal of Morphology, Vol. V.
1890. Tassari, Roméo. Sull' primo di sviluppo dei Teleostei.
Atti della R. Accademia dei Lincei Series 4, Vol. VII.
Resumé de l'Auteur, Archives Italienne de Biologie,
Vol. VIII. 1893.
1890. Grönroos, S. Några af Fiskarna vid den Finska
Kustens.

1888. Fessenden, Felix. Recherches sur le développement des
 Poissons Osseux: Embryologie de la Poisson. Journal
 de l'Anatomie et de Physiologie. 3. VIII.
1893. Herwig, Richard. Reife und Reifwerden, Der Furch-
 ungsprozess, in Verhandl. vergl. u. exper. Entwickel-
 ungslehre der Wirbeltiere. (Fests. von G. Hertwig).
1898. His, Wilhelm. Ueber Furch- und Eneidbildung:
 Studien an Salmonidenkeim. Abh. der anat.-phys.
 Klasse der K. Sächs. Gesellsch. der Wissenschaften,
 Bd. XVII., No. 11.
1898. _____. Protoplasmastrahlen an Salmonidenkeim.
 Idem. Bd. XVII., No. III.
1891. Hoffmann, C. H. Zur Ontogenie der Knochenfische.
 Verhandlungen der Koninklijke Akademiën van Wetenschap-
 pen (Amsterdam) deel XXI.
1896. _____. Ueber den Ursprung und die Bedeutung der
 sogenannten "freien" Kerne in der Keimungsperiode bei
 den Knochenfische. Zeitschrift für Wissensch.
 Zoologie. Bd. XLVI.
1892. Liot, André. Recherches sur les Poissons Lophobranchi-
 es. Annales des Sciences Naturelles. Ser. 3, Vol. VIII.
1891. T. O. The Spermatozooids of Fishes.
 Journal of Morphology, Vol. 1.

1852. Kingsley, T. C. and Tom, H. W. Some observations on the Embryology of the Teleosts. Boston Society of Natural History, Memoirs, Vol. III.
1872. Klein, F. Researches on the First Stages of the Development of the Common Trout. Monthly Microscopical Journal. Vol. VII.
1871. Kopsch, Fr. Die Entstehung des Eitersackepithels und die Bildung der Keimhaut. Internat. Monatschrift für Anat. und Phys. Bd. XVIII.
1888. Kovalevski, M. Ueber die ersten Entwicklungsprozesse der Knochenfische. Zeitschrift für Wissenschaft. Zoologie, Bd. XVIII.
1888. Krojer, Henrik. Danmarks Fiske, Tredie Binde. Hjöbenhavn
1888. Kyffer, N. Beobachtungen uober die Entwicklung der Knochenfische, Archiv für Mikr. Anat. Bd. IV.
1871. LaFont, A. Note pour servir à la Faune de la France, etc. Actes de la Société Linnéenne Française, Tome VIII.
1891. Milljeborg, M. Mykobranchii. Överiges och Förges Fiskar, Tredje Boken. Upsala.
1887. Nist, J. H. Zur Entwicklungsgeschichte der Knochenfische. Zeitschrift für Wissenschaft. Zoologie. Bd. XIV.

1854. Lwoff, P. Die Fische der Provinz Meigsbüttel und die Faunisten der Chocoma-Gruppe des Vordons bei den Fischthiere. Knochenfische. Mémoires Société Impériale des Naturalistes de Moscou, Nouvelle Sér., 3.
1874. Mats, A. S. De den Dreendegede Kathaleis --- Syngnathus typus. -- Utvevelig den Fortplanting. Kongl. Diss. Lund.
1882. Mats, A. Note sur ----- un Syngnathus à deux Queue. Annales des Sciences Naturelles Zool., T. XVIII.
1878. Obelacher, Joseph. Beiträge zur Entwicklungsgeschichte der Knochenfische nach Beobachtungen an Fischfossilien. Zeitschrift für Wiss. Zool., Bd. XIII.
1894. Golovitzza, Paul G. Notes de Biologie -- Appendement et Fécondation chez l'Actopus vulgare Linn. Archiv de Zoologie Experimentale, Ser. 3, T. II.
1867. Ranson, J. Observations on the Ovary of Osseous Fishes. Phil. Trans., Vol. 103.
1838. Retke, Heinrich. Zur Anatomie der Fische. Archiv für Anatomie und Physiologie Bd.
1837. _____ Ueber die Entwicklung der Syngnathen zur Morphologie: Reise-bemerkungen aus Schweden.
1840. _____ Bemerkungen Ueber Syngnathus neptoreus, etc. Archiv für Anatomie und Physiologie, Bd.

1883. Rebecq, A. *Veens Grundläggningar och Samfundss der Färdig*
Morphologiska Samfundet. Bd. VIII.
1883. Reichard, Jacob. *The Rip- Eggs and the Generation*
of the Fall-Steel Pike. Report Michigan Board Fish
Commissioners. Lansing.
1883. Retzius, A. *Anatomisk undersökning öfver äggceller*
af Squamatus 1881 och öpnin. Kongliga Svenska
Vetenskaps, Handlingar. Stockholm.
1882. Snyder, John A. *A Contribution to the Development and*
Morphology of the Tophobranchiates (Tilpocampus hudson
ius). Bulletin U. S. Fish Commission, Vol. I. for 1881.
1884. _____ *A Contribution to the Embryology of*
Ossseous Fishes, Etc., Etc. Report of U. S. Fish
Commission, Part V. for 1882.
1887. _____ *On the Development of Ossseous Fishes, in-*
cluding Marine and Fresh Water Forms. Report U. S.
Fish Commission, Part VIII, for 1886.
1888. Semassa, Paul. *Studien über den Einfluss des Dotters*
auf der Gastrulation und die Bildung der primären
Keimblätter der Wirbelthiere. III. Vertebraten. Archiv
für Entwicklungsmechanik der Organismen. Bd. III.
1885. Stricker, Salomon. *Untersuchungen über die Entwickel-*
ung der Sacnforelle. Sitzber. der K. Akad. der Wissen-
schaften. Wien.

1838. Vagt, P. and Farnstrand. Recherches sur l'Anatomie comparée, etc., etc. Des Femmes des Lapons du Nord. Annales des Sciences Naturelles, Serie IV., Tome VI.
1831. Wilson, Henry J. The Embryology of the Sea Bass (*Cercanus atrarius*). Bulletin U. S. Fish Commission, Vol. IX, for 1898.
1882. Ziegler, F. Die Embryonale Entwicklung von *Salmo salar*. Inaug. Diss. Freiburg.
1887. _____ Die Entstehung des Plutes bei Knochenfischebrute. Archiv für ik. nat. Bd. XXX.
1890. _____ Die Entstehung des Periplutes bei den Knochenfische. Anatomischer Anzeiger, Bd. 17.

Walter Willis Turley was born August 18, 1886, at Nashville, N. C., and received his liberal education in the public and private schools of that town. He was a student at Short and Terry College, Va., 1907-08, and 1908-09.

After attending for three years in public schools, he went to Nashville, Tenn., and in 1904 was graduated with the degree of B. S. from the Peabody Normal College. In 1905 he received the degree of A. B., and in 1906 that of M. A., from the University of Nashville.

He was Assistant Principal of Orange Street High School, Asheville, N. C., 1907-08, and principal for one year and instructor in Asheville Female College. From 1908-1911, he was instructor in Sciences in Tenney High School, Little Rock, Ark. He entered Johns Hopkins University in 1911, and from 1912-14 was student assistant in General Biology. His subjects were Zoology, Botany, and Zoology.

